

Concomitant p53 Mutation and MYCN Amplification in Neuroblastoma

Reinaldo Manhani,¹ Lilian Maria Cristofani, MD, PhD,² Vicente Odone Filho, MD, PhD,²
and Israel Bendit, MD^{1*}

The MYCN oncogene is amplified in 20% of childhood neuroblastoma and is associated independently with poor prognosis. Alteration of the p53 tumor suppressor gene, in contrast, occurs infrequently in these tumors. In this report, we described a 3-year-old girl with stage IV neuroblastoma. Molecular analysis revealed

both MYCN gene amplification and a point mutation of the p53 tumor suppressor gene. To our knowledge, this is the first reported case of neuroblastoma with genetic alterations of both these genes. *Med. Pediatr. Oncol.* 29:206–207, 1997. © 1997 Wiley-Liss, Inc.

Key words: neuroblastoma; p53 mutations; MYCN gene amplification

INTRODUCTION

Neuroblastoma (NB) is the most common extracranial solid tumor of childhood [1]. Studies on MYCN oncogene amplification as a prognostic marker of poor outcome in these neoplasms have served as a paradigm for translational studies [2]. Additional genetic alterations that occur in neuroblastoma and which have also been correlated with poor outcome include LOH for chromosomes 1 and 14, trisomy of 17 q, and low expression of TRK-A [3–11].

Mutation of the p53 tumor suppressor gene on chromosome 17p13 is the most frequent genetic alteration in human cancers to date [12,13]. However, in neuroblastoma, as in many pediatric solid tumors, p53 mutations occur infrequently [14–18]. In this report we describe a case of stage IV NB in which both MYCN amplification and point mutation of the p53 tumor suppressor gene were identified.

MATERIAL AND METHODS

Case Report

A 3-year-old girl presented with right adrenal and periorbital masses, bone pain, pallor, and fever. Bone marrow analysis revealed infiltration with neuroblastoma. Bone lesions were diffuse and CT revealed meningeal involvement. She was treated with intensive chemotherapy (Carboplatin, Etoposide, Cyclophosphamide, Doxorubicin) and surgery. She eventually died of progressive disease.

Molecular Analysis

High molecular weight DNA was extracted from tumor samples obtained at the time of diagnosis and second-look surgery following chemotherapy using proteinase as described elsewhere [19]. MYCN amplification was measured by Southern blot analysis as described

previously [19]. p53 mutations were identified by direct nucleotide sequence analysis of PCR products using Sequenase™ (AMERSHAM International, Buckinghamshire, UK), oligonucleotide primers, and conditions described elsewhere [20].

RESULTS AND DISCUSSION

As part of a protocol designed to determine prognostic markers in NB, MYCN gene amplification in the tumor specimen was determined by Southern blot analysis. The tumor specimen at diagnosis and after chemotherapy contained more than 200 copies of the MYCN oncogene. This finding predicted a poor outcome for the patient, which was unfortunately confirmed when she died 9 months after diagnosis despite intensive chemotherapy.

The tumor specimen from both occasions was also tested for mutation of the p53 tumor suppressor gene by direct DNA sequence analysis of exons 5–8. A G→T transversion was identified (Cys→Phe) at codon 277 that alters the charge of the amino acid (Fig. 1) in the tumor specimen obtained at second-look surgery after chemotherapy; p53 mutations were not detected in the tumor sample obtained at the time of diagnosis (or in the blood from the patient, confirming that this is a *de novo* mutation and not a polymorphism). This exact mutation has not been described before, but similar alterations at this codon, which is a hotspot for mutation, have been de-

¹Research and Molecular Biology Division, Pró-Sangue Hemocentro de São Paulo Foundation, São Paulo, Brazil; ²Children's Institute of the University of São Paulo Medical School, São Paulo, Brazil

*Correspondence to: Israel Bendit, M.D., Research and Molecular Biology Division, Pró-Sangue Hemocentro de São Paulo Foundation Av. Dr. Enéias de Carvalho Aguiar, 155 1st floor CEP 05403-000, São Paulo, Brazil.

Received 22 July 1996; Accepted 5 November 1996




Fig. 1. DNA sequence of p53 exon 8 in DNA from neuroblastoma specimen which exhibited anomalous migration on SSCP analysis (data not shown). p53 exon 8 DNA including codon 276 and 278 from neuroblastoma obtained at second look is at left. Wild-type p53 exon 8 DNA sequence from neuroblastoma at diagnosis is at right.

scribed [21,22]. The p53 mutation, which occurs at a CpG dinucleotide, may have been present but undetected in a small subset of original tumor cells that subsequently expanded, leading to its later identification in the second-look surgical specimen. Alternatively, the mutation could have arisen *de novo* as a result of the chemotherapy. In either case, detection of p53 mutation was associated with a progressive tumor that was unresponsive to chemotherapy. p53 mutations can be associated paradoxically with either increased or decreased sensitivity to chemoradiotherapy [23].

There are several possible explanations for the infrequency of p53 mutation in NB as compared to the prevalence in other common solid tumors in childhood [24–28]. It is possible that other mutations (e.g., deletions on chromosome 1 or 14) affect the pathway.

While other studies confirm the frequency of MYCN amplification and infrequency of p53 mutations in neuroblastoma [14–18], to our knowledge, this is the first case in which both genetic alterations were observed to occur in the same tumor.

REFERENCES

- Hayes FA, Smith EI: Neuroblastoma. In P.A. P, D.G. P, (eds.): "Principles and Practice of Pediatric Oncology." Philadelphia: Lippincott, 1989:607–622.
- Seeger RC, Brodeur GM, Sather H: Association of multiple copies of N-myc oncogene with rapid progression of neuroblastomas. *N Engl J Med* 313:1111–1116, 1985.
- Gilbert F, Feder M, Balaban G, Brangman D, Lurie DK, Podolsky R, Rinaldi V, Vinikoor N, Weisband J: Human neuroblastomas and abnormalities chromosomes 1 and 17. *Cancer Res* 44:5444–5449, 1984.
- Fong CT, Dracopoli NC, White PS: Loss of heterozygosity for the short arm of chromosome 1 in human neuroblastomas: correlation with N-myc amplification. *Proc Natl Acad Sci USA* 86:3753–3757, 1989.
- Martinsson T, Weith A, Cziepluch C, Schawab M: Chromosome 1 deletion in human neuroblastomas: generation and fine mapping of microclones from the distal 1 p region. *Genes Chromosomes Cancer* 1:67–78, 1989.
- Fong CT, White PS, Sapienza C: Loss of heterozygosity for chromosomes 1 or 14 defines subsets of advanced neuroblastomas. *Cancer Res* 52:1780–1785, 1992.
- Takayama H, Suzuki T, Mugishima H: Deletion mapping of chromosomes 14q and 1p in human neuroblastoma. *Oncogene* 7:1185–1189, 1992.
- Borrello MG, Bongarzone I, Pierotti MA: *trk* and *ret* proto-oncogene expression in human neuroblastoma specimens: High frequency of *trk* expression in non-advanced stages. *Int J Cancer* 54:540–545, 1993.
- Nakagawara A, Arima-Nakagawara M, Scavarda NJ, Azar CG, Cantor AB, Brodeur GM: Association between high levels of expression of the *trk* gene and favorable outcome in human neuroblastoma. *N Engl J Med* 328:847–854, 1993.
- Caron HN: Allelic loss of chromosome 1 and additional chromosome 17 material are both unfavorable prognostic markers in neuroblastoma. *Med Pediatr Oncol* 24:215–221, 1995.
- Caron H, van Sluis P, de Kraker J: Allelic loss of chromosome 1p as a predictor of unfavorable outcome in patients with neuroblastoma. *N Engl J Med* 334:225–230, 1996.
- Levine AJ, Momand J, Finlay CA: The p53 tumor suppressor gene. *Nature (Lond)* 351:453–456, 1991.
- Hollstein M, Sidransky D, Vogelstein B, Harris CC: p53 mutations in human cancers. *Science* 253:49–53, 1991.
- Imamura J, Bartran CR, Berthold F, Harms D, Nakamura H, Koefler HP: Mutation of the p53 gene in neuroblastoma and its relationship with N-myc amplification. *Cancer Res* 53:4053–4058, 1993.
- Komuro H, Hayashi Y, Kawamura M: Mutations of the p53 gene are involved in Ewing's sarcoma but not in neuroblastoma. *Cancer Res* 53:5284–5288, 1993.
- Vogan K, Bernstein M, Leclerc JM: Absence of p53 gene mutations in primary neuroblastomas. *Cancer Res* 53:5269–5273, 1993.
- Castresana JS, Bello MJ, Rey JA: No TP53 mutations in neuroblastomas detected by PCR-SSCP analysis. *Genes Chromosomes Cancer* 10:136–138, 1994.
- Hosoi G, Hara J, Okamura T: Low frequency of the p53 gene mutations in neuroblastoma. *Cancer* 73:3087–3093, 1994.
- Garvin JJ, Bendit I, Nisen PD: N-myc expression and amplification in metastatic lesions of stage IV-S neuroblastoma. *Cancer* 65:2572–2575, 1990.
- D'Amico D, Carbone D, Mitsudomi T: High frequency of somatically acquired p53 mutations in small cell lung carcinoma cell lines and tumors. *Oncogene* 7:339–346, 1992.
- Kovar H, Auinger A, Jug G: Narrow spectrum of infrequent p53 mutations and absence of MDM2 amplification in Ewing tumours. *Oncogene* 8:2683–2690, 1993.
- Seth A, Palli D, Mariano JM: p53 gene mutations in women with breast cancer and a previous history of benign breast disease. *Eur J Cancer* 30A:808–812, 1994.
- Russel SJ, Yun WY, Waber PG, Shuford M, Schold C, Nisen PD: p53 mutations, O⁶-alkylguanine DNA alkyltransferase activity, and sensitivity to procarbazine in human brain tumors. *Cancer* 75:1339–1342, 1995.
- Toguchida J, Yamaguchi T, Dayton SH: Prevalence and spectrum of germline mutations of the p53 gene among patients with sarcoma. *N Engl J Med* 326:1301–1308, 1992.
- von Deimling A, Eibl RH, Ohgaki H: p53 mutations are associated with 17p allelic loss in grade II and grade III astrocytoma. *Cancer Res* 52:2987–2990, 1992.
- Frankel RH, Bayona W, Koslow M, Newcomb EW: p53 mutations in human malignant gliomas: Comparison of loss of heterozygosity with mutation frequency. *Cancer Res* 52:1427–1433, 1992.